

APPROXIMATION AND SPANNING IN THE HARDY SPACE, BY AFFINE
SYSTEMS

H.-Q. Bui¹ and **R.S. Laugesen**²

¹ *Department of Mathematics & Statistics,
University of Canterbury,
Private Bag 4800, Christchurch, New Zealand.*

² *Department of Mathematics,
University of Illinois,
Urbana, IL 61801, U.S.A.*

Report Number: UCDMS2005/6

July 2005

Keywords: Completeness, quasi-interpolation, approximate identity,
atomic, scale averaging.

APPROXIMATION AND SPANNING IN THE HARDY SPACE, BY AFFINE SYSTEMS

H.-Q. BUI AND R. S. LAUGESEN

ABSTRACT. We find weak conditions on $\psi \in L^1(\mathbb{R}^d)$ with $\widehat{\psi}(0) = 1$ such that every function in the Hardy space is a linear combination of translates and dilates of ψ . More precisely, we prove for each $f \in H^1(\mathbb{R}^d)$ the scale averaged approximation formula

$$f(x) = \lim_{J \rightarrow \infty} \frac{1}{J} \sum_{j=1}^J \sum_{k \in \mathbb{Z}^d} c_{j,k} \psi(a_j x - k) \quad \text{in } H^1(\mathbb{R}^d),$$

where $\{a_j\}$ is an arbitrary lacunary sequence (such as $a_j = 2^j$) and the coefficients $c_{j,k}$ are local averages of f . This holds in particular if ψ is Schwartz class, or if $\psi \in L^p$ (for some $1 < p < \infty$) has compact support. A corollary is a new affine decomposition of H^1 in terms of differences of ψ .

—

Date: July 7, 2005.

1991 Mathematics Subject Classification. Primary 42A50, 42B30, 42C30.

Key words and phrases. Completeness, quasi-interpolation, approximate identity, atomic, scale averaging.

Laugesen was partially supported by N.S.F. Award DMS-0140481, a Maclaurin Fellowship from the New Zealand Institute of Mathematics and its Applications, and a Visiting Erskine Fellowship from the University of Canterbury.